

Integration of Transcriptional and Mutational Data improves the Stratification of

Peripheral T-Cell Lymphoma

Francesco Maura^{1-2*}, Luca Agnelli^{2-3*#}, Daniel Leongamornlert¹, Niccolò Bolli^{1-2,5}, John Chan⁴, Anna Doderio⁵, Cristiana Carniti⁵, Tayla Heavican⁶, Alessio Pellegrinelli⁷, Giancarlo Pruneri⁷, Adam Butler¹, Shriram Bhosle¹, Annalisa Chiappella⁸, Alice Di Rocco⁹, Pier Luigi Zinzani¹⁰, Francesco Zaja¹¹, Roberto Piva¹², Giorgio Inghirami^{12,13}, Wenyi Wang¹⁴, Teresa Palomero¹⁵, Javeed Iqbal⁶, Antonino Neri²⁻³, Peter J Campbell¹ and Paolo Corradini^{2,5#}

¹Cancer, Ageing and Somatic Mutation Programme, Wellcome Trust Sanger Institute, Hinxton, United Kingdom; ²Department of Oncology and HematoOncology, University of Milan, Milan, Italy; ³Hematology, Fondazione IRCCS Ca' Granda Ospedale Maggiore

Policlinico, Milan, Italy;

⁴City of Hope, Duarte, CA;

⁵Department of Hematology, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy;

⁶Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha;

⁷Department of Pathology and Laboratory Medicine, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy;

⁸Department of Hematology, Azienda Ospedaliera Città della Salute e della Scienza, Torino, Italy;

⁹Sapienza University of Rome, Rome, Italy;

¹⁰Institute of Hematology, University of Bologna, Bologna, Italy

¹¹Clinica Ematologica, DAME, University of Udine, Udine, Italy;

¹²Center for Experimental Research and Medical Studies, University of Torino, Department of Molecular Biotechnology and Health Sciences, Torino, Italy;

¹³Pathology and Laboratory Medicines, Weill Cornell Medical College,

¹⁴The University of Texas MD Anderson Cancer Center, Houston

¹⁵Institute for Cancer Genetics, Columbia University, New York, NY 10032

*The first two authors equally contributed

Key words: PTCL, Gene expression profile, CIBERSORT, next-generation sequencing

Corresponding Author:

Luca Agnelli

Department of Oncology and Hemato-Oncology, University of Milan, Milan, Italy

F. Sforza, 35, 20122 Milan Italy

luca.agnelli@unimi.it

Tel (+39)0255033328

Fax: (+39)0255034571

Abstract

The histological diagnosis of peripheral T-cell lymphoma (PTCL) can represent a challenge, particularly in the case of close related entities like angio-immunoblastic T-lymphomas (AITL) and PTCL-not otherwise specified (PTCL-NOS). Although gene expression profiling and next generations sequencing have been proven to define specific features recurrently associated with distinct entity, genomic-based stratifications have not yet led to definitive diagnostic criteria and/or entered into the routine clinical practice.

Herein, to improve the current molecular classification between AITL and PTCL-NOS, we analyzed the transcriptional profiles from 503 PTCLs stratified according to their molecular configuration and integrated them with genomic data of recurrently mutated genes (*RHOA*^{G17V}, *TET2*, *IDH2*^{R172}, and *DNMT3A*) in 53 cases (39 AITLs and 14 PTCL-NOSs) included in the series. Our analysis unraveled that the mutational status of *RHOA*, *TET2* and *DNMT3A* poorly correlated, individually, with peculiar transcriptional fingerprint. Conversely, a strong transcriptional signature was identified in *IDH2*^{R172} samples. The integrated analysis of clinical, mutational and molecular data led to 19-genes model whose expression can more precisely differentiate the main PTCL nodal entities. According to the new proposed stratification, *RHOA*^{G17V}, *TET2* and *IDH2*^{R217} mutations emerged as more robust diagnostic markers than previously reported for the differentiation between AITL and PTCL-NOS.

This new gene classifier is simple and reproducible and able to improve the current molecular classification and diagnosis of PTCL.

Introduction

Peripheral T-cell lymphomas (PTCL) represent a heterogeneous group of nodal and extra-nodal mature T-cell neoplasms accounting for approximately 10-15% of all lymphoma in the Western countries.¹⁻⁵ The current WHO classification recognizes several distinct subsets, and multiple provisional entities have been recently proposed.³ The most frequent categories include PTCL-not otherwise specified (PTCL-NOS), angioimmunoblastic lymphoma (AITL), and anaplastic large T cell lymphoma (ALCL) with or without translocations involving *ALK*.³ Overall, these entities encompass approximately 60% of all PTCL. With the exception of ALCL with *ALK* translocations, PTCL have an aggressive clinical course and poor response to conventional chemotherapy. Although new agents have been recently approved and several others are in clinical trials, the overall clinical success remains unsatisfactory.^{1,6-8} This is partially due to a limited knowledge of PTCL biology and reproducible/informative pre-clinical models, which have significantly hindered any mechanistic study aimed at testing candidate targeted treatments.

Even though pathological diagnoses can be rendered in many patients, difficulties can occur, and in particular for those samples sharing features borderline between AITL and PTCL-NOS.⁹ Previous studies have shown that these two latter entities might bear distinct gene expression profiles (GEP).¹⁰⁻¹⁶ However, clinical usage of GEP has been limited due to its technical availability and to the absence of a consensus gene signature. Recent advances of next generation sequencing (NGS), and the discovery of recurrent mutated genes (*RHOA*, *TET2*, *DNMT3A*) in approximately 60-70% of AITL and in 20-30% of PTCL-NOS, have somehow changed this landscape.^{15,17-20} Notably, 20-30% of AITLs cases can carry hotspot *IDH2*^{R172} mutations that are virtually absent in PTCL-NOSs.¹⁷ Nevertheless, these findings have not yet significantly impacted the clinical daily practice, which largely relies on histology and phenotype of tumor cells.³ Moreover, while some mutations appear to be linked to distinct GEP signature(s),¹⁵ the full potential of an integrated genotypic-transcriptomic analysis has not been thoroughly tested.

Here, we collected a large GEP dataset of PTCL, and performed an integrative analysis with mutational data to improve the accuracy of disease classification, aiming of recognizing and more precisely stratifying AITL/PTCL-NOS lymphoma.

Material and Methods

We analysed 503 PTCL and 38 normal T-cell cases, univocally acquired from 8 studies (GSE6338, GSE14879, GSE19067, GSE19069, GSE58445 and GSE65823 at <http://www.ncbi.nlm.nih.gov/geo/>; ETABM702 and ETABM783 at <https://www.ebi.ac.uk/arrayexpress>) (**Supplementary Figure 1**).^{10,12,21-24} Normalized data were extracted from CEL files using RMA procedure and the annotation for HG-U133Plus2.0 arrays available at

<http://brainarray.mbni.med.umich.edu/Brainarray/Database/CustomCDF/21.0.0/entrezg.asp>.

A batch-effect correction was applied as previously described.^{25,26} The whole data set with all available clinical and genomic information acquired was uploaded to <https://github.com/emacgene/PTCL>.

To investigate the contribution of recurrent mutations to the transcriptional pattern, we adapted a recently published analysis²⁷ to 39 AITLs and 14 PTCL-NOSs for whom mutational data for *IDH2*^{R172}, *DNMT3A*, *TET2* and *RHOA* were available.¹⁵ Next, we created a model that included the lymphoma histotype, gene expression profile, genotype, and age and gender of each patient. Finally, *ConsensusClusterPlus* package for R²⁸ was used to determine the significance and robustness of natural grouping of patients based on selected transcriptional data, using Ward and Euclidean as linkage and distance metrics, respectively. CIBERSORT analysis was performed as previously described, using standard procedure and LM22 signature.²⁹ Contingency analyses were performed by Fisher's exact test (*fisher.test* R function). The CIBERSORT different contribution for each signature was then tested by *pairwise.wilcox.test* R function. Benjamini-Hochberg correction was used for multiple testing adjustment. The pathway enrichment analysis was performed using different modalities. First, the *tmod* R package was used on *limma*-derived signatures to decipher

whether clusters deregulate blood cell-associated transcriptional modules described by
Chaussabel *et al.*³⁰ and by Li *et al.*,³¹ according to the procedure described by Weiner et al.³²
The full analysis process written in R is provided in **Supplementary Data 1**. Second, gene
set enrichment analysis was run on extended gene set modules using the GSEA java tool
from the Broad institute (<http://software.broadinstitute.org/gsea/index.jsp>). In this second
analysis, to reduce the incidence of false positives results, we choose the *samr* package,
and a more conservative approach, to list differentially expressed genes.

To validate our findings in an independent dataset, we imported data from 34
previously published RNAseq samples (dbGap accession n. phs000689.v1.p1).¹⁷ The
RNAseq raw expression data were normalized as previously described.³³The full
computational process written in R is provided in **Supplementary Data 2**.

Results

Dataset definition

We analysed the transcriptional profiles of 503 PTCL cases and 38 normal donors provided by 8 different studies.^{10,12,13,21-24,34} According to the most recent updates in the T-cell lymphoma molecular classification,³ these series included the following histotypes: 127 AITL, 144 PTCL-NOS, 56 *ALK*+ ALCL, 96 *ALK*- ALCL, 21 Adult T-Cell Lymphoma (ATLL), 59 NK/T-cell lymphomas (**Figure 1a**). By unsupervised hierarchical clustering and principal component analysis, *ALK*- and *ALK*+ ALCLs, ENKTL and ATLL were associated with distinct signatures; conversely, the transcriptional portrait of AITL and PTCL-NOS displayed a considerable overlap (**Figure 1b-c**).

IDH2^{R172} but not *RHOA*, *TET2* and *DNMT3A* is associated with specific transcriptional pattern

To improve the distinction between AITL and PTCL-NOS, we integrated expression profiles with clinical, laboratory and mutational data (39 AITLs and 14 PTCL-NOSs) (See **Material and Methods** and **Supplementary Data 1**). This strategy lead to a preliminary list of 221/20.363 (1%) differentially modulated genes [false rate discovery (FDR) <1%] (**Figure 2a** and **Supplementary Table 1**). Among these, 30 of them emerged as significantly and specifically associated to one distinct variable. Gender selectively impacted 14 genes located on the X and Y chromosome and for this reason they were not included for further analysis. Interestingly, the mutational status of *RHOA*^{G17V}, *TET2* and *DNMT3A* poorly correlated with any distinct transcriptional profile in both AITL and PTCL-NOS, as no gene showed differential expression to be correlated with the genotype (**Figure 2b**). Conversely, *IDH2*^{R172} samples were linked to a unique transcriptional signature including 3 genes with FDR <1%: *ID2*, *NETO2* and *SLC5A3* (**Supplementary Data 1** and **Figure 2b**). Notably, AITL and PTCL-NOS transcriptional profiles could be differentiated by the expression of 13 genes, including 3 of them recently described in recent gene expression model (*ROBO1*, *ARHGEF10* and *EFNB2*) (**Figure 2a-b**).¹²

Lastly, we decided to combine the signature specifically associated with *IDH2*^{R172} mutational status (n=3) and those linked to histological diagnoses (n=13) to all AITL and PTCL-NOS (n=271) samples. This approach led to a hierarchical clustering with an improved clustering of PTCL-NOS and AITL cases. Collectively, these data are consistent with biologically distinct subgroups associated with specific mutational-based signatures, and linked to defined pathological features (**Figure 2c**).

A 19-gene expression-based model distinguished PTCL subtypes

Our group has previously described a 3-gene signature capable to effectively distinguish *ALK*-ALCLs from PTCL-NOS³⁵. Here, we combined this 3-gene signature with our new 16-gene model, ultimately leading to a 19-gene model, aimed at improving the stratification of all major PTCL entities (*ALK*-ALCL, PTCL-NOS and AITL). This led to the recognition of five distinct subsets (**Figure 3, Supplementary Data 1 and Table 1**). The first group (C-1; n=87) was mainly composed by AITL samples (93%; 81/87), and enriched by *IDH2*^{R172} (16/36; 44%) and *RHOA*^{G17V} (23/29; 79%) mutated lymphoma. The second (C-2; n=103), with an AITL-like signature, included samples annotated as well as PTCL-NOSs (64/103; 62%) and AITLs (33/103; 32%). In this group, *IDH2*^{R172} samples were minimally represented compared to C-1 (1/23; p=0.001); conversely, a higher prevalence of *RHOA*^{G17V} mutations was observed (15/29; 52%); particularly higher than that seen in cluster 4 (p=0.005). The third cluster represented only by 21 patients, included samples that could be linked to any of 3 main PTCL entities. Group 4 (C-4; n=63) included largely PTCL-NOSs (52/63; 82%), displaying a distinctive transcriptional profile characterized by a low prevalence of *RHOA* and *TET2* mutations compared to C-1 (p<0.0001 and p=0.0002 respectively) or C-2 (p=0.005 and 0.03 respectively) (**Supplementary Figure 2**).¹² In the last cluster (C-5; n=55) *ALK*-ALCL cases (47/69; 68%) were over-represented, confirming the strong association between these lymphoma and the expression of *TNFRSF8*, *BATF3* and *TMOD1*.³⁵ No significant differences were observed in the *DNMT3A* mutational profile between any cluster (**Table 1**). Overall, this approach could improve the previous molecular

characterization of up to 8% AITLs, 31% *ALK*- ALCL and 29% PTCL-NOSs providing the bases of the recognition of unique biological subgroups (**Table 1**).

Next, to discover potential gene classifiers, we investigated the global gene expression among samples within the 5 clusters (**Supplementary Data 1**). Firstly, we confirmed molecular signatures previously associated to AITL and PTCL-NOS (i.e *ROBO1*, *LPAR1*, *SOX8*, *TUBB2B*, *TNFRSF8*, *TMOD1*, *BATF3*, *ATP6V0D1*, *CHI3L1*, *CREG1*, *CTSB*, *CTCS*, *FTL*, *HCK*), strengthening the ability of the 19-gene model to differentiate distinct pathological entities.^{12,36}

Running 2 independent pathway analyses (**Material and Methods and Supplementary Data 1**), we observed a significant enrichment of B-cell and Plasma cell pathways among C-1 and C-2, in agreement with the notion that AITL are often enriched in B-cells/plasma cells; conversely the cluster 5 showed an enrichment for T-CD4 cell cycles pathway, but a significant low involvement of ITK/PKC and T-cell activation pathway (**Supplementary Figure 3**), in agreement with the consideration that the TCR signalling has been reported to be non-dominant in this group.³⁷

Validation Cohort

To validate our 19-gene signature, we analyzed the RNAseq expression of 34 PTCL-NOSs (11 AITL, 11 PTCL-NOS, 8 *ALK*- ALCL and 4 *ALK*+ ALCL) from a previously published dataset.¹⁷ The expression data of 2 of 19 genes were not available (*AL441992.1* and *SLC5A3*). Using the remaining 17 genes we were able to extract 4 main clusters (**Supplementary Data 2 and Figure 4**). The first cluster was compatible with the C-1 and C-2 (AITL and AITL-like) clusters. All *RHOA*, *TET2* and *IDH2*^{R172} mutated cases were included in this cluster mostly composed by a fraction of PTCL-NOSs (6/11) and all AITLs (10/10). The second was characterized by a profile similar to the C-4 group, with PTCL-NOS without *RHOA* and *TET2* mutations. The remaining clusters included the great majority of *ALK*+ and *ALK*- ALCL cases (9/11).

Computational investigation defines distinct tumor-host compositions

Seeking to discover whether the new PTCL clusters may be linked/associated to any host-tumor features, we used CIBERSORT to define the cellular composition of the tumor and host elements. CIBERSORT is a computational tool that accurately resolves relative fractions of different elements within complex environments (**Figure 5a** and **Supplementary Data 1**).²⁹ This analysis revealed that the C-1 and C-2 clusters were characterized by a highest contribution of plasma cell and B-cell signatures (**Figure 5b**). Conversely, both clusters displayed a very low T-reg but high macrophages M₀ signals. Cluster 5, mainly represented by ALK- ALCLs, was enriched for activated T-cell CD4+ and by Macrophages (M₁ and M₂) signatures (**Figure 5b** and **Supplementary Data 1**). Notably, the contribution of T-follicular helper cells was significantly lower in C5 than in the C-1, C-2 and C-4 groups. Lastly, the C-3 cluster was characterized by a unique profile associated with NK-cell (p<0.01) and activated mast cell (p<0.001) signatures, likely reflecting unique features of these lymphoma compared to those seen on bona fide AITL, PTCL-NOS and ALCL cases (**Supplementary Data 1**).

Discussion

GEP has emerged as one of the most robust and reliable approach to differentiate human lymphoma. This has also been true for T-cell lymphoma, a strategy that have allowed the identification of even closely related entities like AITL and PTCL-NOS.^{10,12,22-24,34,35} However, GEP profile from formalin fixed paraffin embedded samples has been technical problematic and thus the molecular/expression stratification of PTCL has not entered yet into the routine clinical practice. Recently, novel technologies (e.g. NanoString) have provided for the first time a reproducible and feasible quantification of specific transcripts, mainly for diffuse large B-cell lymphoma.³⁸⁻⁴⁰ conversely, this approach has not yet been tested in PTCL, likely because a short and robust list of differentially expressed genes has not emerged yet.

To bridge this gap, we performed meta-analysis of the largest PTCL series collected to date to assess the molecular profile of the main PTCL subgroups and define a usable list of significant differentially expressed genes. By integrating NGS and GEP data, we were able to discover the transcriptional impact of the recurrent mutations of AITLs and PTCL-NOSs. Indeed, we found that *TET2*, *DNMT3A*, *RHOA*^{G17V} mutations did not show any distinct gene expression signatures. This last mutation was detected at sub-clonal level in a significant fraction of these cases and this may explain its low impact on transcriptional configuration.¹⁵ *TET2* and *DNMT3A* mutations are likely as early as hematopoietic stem cells mutations involved in clonal hemopoiesis;^{41,42} therefore, we could not exclude that *TET2*-associated gene expression signature might be shared by other subtypes and masked by AITL molecular classification, subsequently limiting its extraction through our statistical process. Conversely, in line with a previous report,¹⁵ the *IDH2*^{R172} mutation significantly correlated with a distinct expression signature independent from the molecular subgroup. By integrating pathology, NGS and GEP analyses we defined a short list of highly predictive genes (n=19) whose differential expression divided the samples into 5 clusters, strongly associated with distinct PTCL entities. Specifically, C-1 and C-2 were characterized by most of the AITL hallmarks. In addition, despite their limited gene expression impact, the great

majority of *RHOA*^{G17V} and *TET2* mutations were grouped together, confirming their potential utility in the diagnostic process of these lymphoma. A significant fraction of PTCL-NOS were included in these 2 AITL clusters sharing high prevalence of *RHOA* and *TET2* mutations and expressing some distinct, and in part previously reported, AITL-associated genes, suggesting a better ability of this model to re-assign more correctly the T-follicular helper PTCLs.

Furthermore, even without a direct GEP impact, *RHOA* mutations were mostly clustered within AITL clusters, emerging together with *IDH2*^{R217} mutations as a potential specific diagnostic marker to be further employed in routine diagnostics. Lastly, The majority of PTCL-NOS not assigned to C-2, composed a unique and distinct cluster (C4).

Based both on the present and recently published^{10,12,37} data, PTCL-NOS group within the C-4 could not be considered any more like an “orphan” or “not-otherwise specified” subset, but instead as a distinct biologically entity. In fact, the lymphoma patients within this cluster have a distinct genotypic and transcriptomic pattern, mostly characterized by a significant enrichment in downregulated genes, confirming previous findings.¹³ Clearly, further studies will be needed to confirm our findings, and to further refine the list of genomic driver events and genes whose expression levels may have diagnostic value. However, our data were preliminarily cross-validated in an independent small RNAseq cohort¹⁷, confirming the robustness of the signature. This was true even across gene expression platforms, and thus represents a promising test bed for future routine nanostring approaches.

Combining the expression levels and the mutational status of a limited set of genes in future studies, we will be able to improve the current diagnostic approach of PTCLs and lay the basis for effective treatments through the identification of recurrently dysregulated pathways.

281 **Acknowledgment**

282 NB is funded by AIRC (Associazione Italiana per la Ricerca sul Cancro) through a MFAG
283 (n.17658). FM is supported by A.I.L. (Associazione Italiana Contro le Leucemie-Linfomi e
284 Mieloma ONLUS)

285

286 **Authorship Contributions:**

287 F.M., L.A. and P.C.: designed the study, collected and analysed the data, and wrote the
288 paper;; D.L., W.W., P.J.C., and S.B. analysed the data.; J.C., T.H., A.D., C.C., A.P., G.P.,
289 A.B., A.C., A.D.R., P.L.Z., F.Z., U.V., R.P., J.I., T.P. and J.I. collected the data; N.B., A.N.,
290 and J.I. critically revised the paper

291

292 **Disclosure of Conflicts of Interest:**

293 No conflict of interests to declare

294

Figure Legends:

Figure 1. a) Molecular composition of the gene expression cohort (541 cases).

Unsupervised hierarchical clustering (b) and PCA analysis (c) on the entire series.

Figure 2. a) Distribution of the variance of expression levels across genes explained by clinical, molecular and genetic alterations (f-test; FDR<1%; n=221). b) Statistically significant mutation expression interaction terms (f-test; FDR<1%), for each alteration and clinical variable. The associated logarithmic expression fold change is indicated by colour. c) Hierarchical clustering of 271 PTCL-NOSs and AITLs based on the 16-significantly extracted genes. The mutational status for *TET2*, *RHOA*, *DNMT3A*, *IDH2*^{R172} was reported on the top.

Figure 3. Heatmap of the 19-genes model including all PTCL-NOS, AITL and *ALK*-ALCL cases (n=367), stratified according to the cluster determined by the *ConsensusClusterPlus* R function.

Figure 4. a) CIBERSORT profile of each enrolled tumour sample (n=503). b) Boxplot of the relative estimated percentage for the most relevant and significant cell types, stratified by groups (Wilcoxon rank sum test for comparisons).

Figure 5. Validation of the 19-gene model on a RNAseq external series.

317 **Supplementary Figure Legend**

318

319 **Supplementary Figure 1.** Summary of the histological composition of each external gene
320 expression data set included in this study.

321

322 **Supplementary Figure 2.** Heat map based on the differential expression of the 16-genes
323 between the C-2 and C-4 PTCL-NOSs.

324

325 **Supplementary Figure 3.** The *tmod* most significant differentially enriched pathway among
326 the extracted 5 clusters,

327

References

- 1 Ellin, F, Landstrom J, Jerkeman M & Relander T. Real-world data on prognostic factors and treatment in peripheral T-cell lymphomas: a study from the Swedish Lymphoma Registry. *Blood* 2014; **124**:1570-1577.
- 2 Maura F, DA, Carniti C, Bolli N. Biology of Peripheral T cell lymphomas - not otherwise specified: is something finally happening? *Pathogenesis* 2016; **3** 9–18.
- 3 Swerdlow, SH, Campo E, Pileri SA, Harris NL, Stein H, Siebert R *et al*. The 2016 revision of the World Health Organization (WHO) classification of lymphoid neoplasms. *Blood* 2016.
- 4 Van Arnem, JS, Lim MS & Elenitoba-Johnson KSJ. Novel insights into the pathogenesis of T-cell lymphomas. *Blood* 2018; **131**:2320-2330.
- 5 Vose, J, Armitage J, Weisenburger D & International TCLP. International peripheral T-cell and natural killer/T-cell lymphoma study: pathology findings and clinical outcomes. *J Clin Oncol* 2008; **26**:4124-4130.
- 6 Corradini, P, Vitolo U, Rambaldi A, Miceli R, Patriarca F, Gallamini A *et al*. Intensified chemo-immunotherapy with or without stem cell transplantation in newly diagnosed patients with peripheral T-cell lymphoma. *Leukemia* 2014; **28**:1885-1891.
- 7 d'Amore, F, Relander T, Lauritzsen GF, Jantunen E, Hagberg H, Anderson H *et al*. Up-front autologous stem-cell transplantation in peripheral T-cell lymphoma: NLG-T-01. *J Clin Oncol* 2012; **30**:3093-3099.
- 8 Gallamini, A, Stelitano C, Calvi R, Bellei M, Mattei D, Vitolo U *et al*. Peripheral T-cell lymphoma unspecified (PTCL-U): a new prognostic model from a retrospective multicentric clinical study. *Blood* 2004; **103**:2474-2479.
- 9 Herrera, AF, Crosby-Thompson A, Friedberg JW, Abel GA, Czuczman MS, Gordon LI *et al*. Comparison of referring and final pathology for patients with T-cell lymphoma in the National Comprehensive Cancer Network. *Cancer* 2014; **120**:1993-1999.
- 10 Iqbal, J, Weisenburger DD, Greiner TC, Vose JM, McKeithan T, Kucuk C *et al*. Molecular signatures to improve diagnosis in peripheral T-cell lymphoma and prognostication in angioimmunoblastic T-cell lymphoma. *Blood* 2010; **115**:1026-1036.
- 11 Iqbal, J, Wilcox R, Naushad H, Rohr J, Heavican TB, Wang C *et al*. Genomic signatures in T-cell lymphoma: How can these improve precision in diagnosis and inform prognosis? *Blood Rev* 2015.
- 12 Iqbal, J, Wright G, Wang C, Rosenwald A, Gascoyne RD, Weisenburger DD *et al*. Gene expression signatures delineate biological and prognostic subgroups in peripheral T-cell lymphoma. *Blood* 2014; **123**:2915-2923.
- 13 Piccaluga, PP, Fuligni F, De Leo A, Bertuzzi C, Rossi M, Bacci F *et al*. Molecular profiling improves classification and prognostication of nodal peripheral T-cell lymphomas: results of a phase III diagnostic accuracy study. *J Clin Oncol* 2013; **31**:3019-3025.

369 14 Piva, R, Agnelli L, Pellegrino E, Todoerti K, Grosso V, Tamagno I *et al.* Gene
370 expression profiling uncovers molecular classifiers for the recognition of anaplastic
371 large-cell lymphoma within peripheral T-cell neoplasms. *J Clin Oncol* 2010; **28**:1583-
372 1590.

373 15 Wang, C, McKeithan TW, Gong Q, Zhang W, Bouska A, Rosenwald A *et al.*
374 IDH2R172 mutations define a unique subgroup of patients with angioimmunoblastic
375 T-cell lymphoma. *Blood* 2015; **126**:1741-1752.

376 16 Wang, T, Feldman AL, Wada DA, Lu Y, Polk A, Briski R *et al.* GATA-3 expression
377 identifies a high-risk subset of PTCL, NOS with distinct molecular and clinical
378 features. *Blood* 2014; **123**:3007-3015.

379 17 Palomero, T, Couronne L, Khiabani H, Kim MY, Ambesi-Impiombato A, Perez-
380 Garcia A *et al.* Recurrent mutations in epigenetic regulators, RHOA and FYN kinase
381 in peripheral T cell lymphomas. *Nat Genet* 2014; **46**:166-170.

382 18 Sakata-Yanagimoto, M, Enami T, Yoshida K, Shiraishi Y, Ishii R, Miyake Y *et al.*
383 Somatic RHOA mutation in angioimmunoblastic T cell lymphoma. *Nat Genet* 2014;
384 **46**:171-175.

385 19 Schatz, JH, Horwitz SM, Teruya-Feldstein J, Lunning MA, Viale A, Huberman K *et al.*
386 Targeted mutational profiling of peripheral T-cell lymphoma not otherwise specified
387 highlights new mechanisms in a heterogeneous pathogenesis. *Leukemia* 2015;
388 **29**:237-241.

389 20 Yoo, HY, Sung MK, Lee SH, Kim S, Lee H, Park S *et al.* A recurrent inactivating
390 mutation in RHOA GTPase in angioimmunoblastic T cell lymphoma. *Nat Genet* 2014;
391 **46**:371-375.

392 21 Eckerle, S, Brune V, Doring C, Tiacci E, Bohle V, Sundstrom C *et al.* Gene
393 expression profiling of isolated tumour cells from anaplastic large cell lymphomas:
394 insights into its cellular origin, pathogenesis and relation to Hodgkin lymphoma.
395 *Leukemia* 2009; **23**:2129-2138.

396 22 Iqbal, J, Weisenburger DD, Chowdhury A, Tsai MY, Srivastava G, Greiner TC *et al.*
397 Natural killer cell lymphoma shares strikingly similar molecular features with a group
398 of non-hepatosplenic gammadelta T-cell lymphoma and is highly sensitive to a novel
399 aurora kinase A inhibitor in vitro. *Leukemia* 2011; **25**:348-358.

400 23 Piccaluga, PP, Agostinelli C, Califano A, Carbone A, Fantoni L, Ferrari S *et al.* Gene
401 expression analysis of angioimmunoblastic lymphoma indicates derivation from T
402 follicular helper cells and vascular endothelial growth factor deregulation. *Cancer*
403 *Res* 2007; **67**:10703-10710.

404 24 Scarfo, I, Pellegrino E, Mereu E, Kwee I, Agnelli L, Bergaggio E *et al.* Identification of
405 a new subclass of ALK-negative ALCL expressing aberrant levels of ERBB4
406 transcripts. *Blood* 2016; **127**:221-232.

407 25 Johnson, WE, Li C & Rabinovic A. Adjusting batch effects in microarray expression
408 data using empirical Bayes methods. *Biostatistics* 2007; **8**:118-127.

409 26 Leek, JT, Johnson WE, Parker HS, Jaffe AE & Storey JD. The sva package for
410 removing batch effects and other unwanted variation in high-throughput experiments.
411 *Bioinformatics* 2012; **28**:882-883.

412 27 Gerstung, M, Pellagatti A, Malcovati L, Giagounidis A, Porta MG, Jadersten M *et al.*
413 Combining gene mutation with gene expression data improves outcome prediction in
414 myelodysplastic syndromes. *Nat Commun* 2015; **6**:5901.

415 28 Wilkerson, MD & Hayes DN. ConsensusClusterPlus: a class discovery tool with
416 confidence assessments and item tracking. *Bioinformatics* 2010; **26**:1572-1573.

417 29 Gentles, AJ, Newman AM, Liu CL, Bratman SV, Feng W, Kim D *et al.* The prognostic
418 landscape of genes and infiltrating immune cells across human cancers. *Nat Med*
419 2015; **21**:938-945.

420 30 Chaussabel, D, Quinn C, Shen J, Patel P, Glaser C, Baldwin N *et al.* A modular
421 analysis framework for blood genomics studies: application to systemic lupus
422 erythematosus. *Immunity* 2008; **29**:150-164.

423 31 Li, S, Roupheal N, Duraisingham S, Romero-Steiner S, Presnell S, Davis C *et al.*
424 Molecular signatures of antibody responses derived from a systems biology study of
425 five human vaccines. *Nat Immunol* 2014; **15**:195-204.

426 32 Weiner, J, 3rd, Parida SK, Maertzdorf J, Black GF, Repsilber D, Telaar A *et al.*
427 Biomarkers of inflammation, immunosuppression and stress with active disease are
428 revealed by metabolomic profiling of tuberculosis patients. *PLoS One* 2012;
429 **7**:e40221.

430 33 Love, MI, Anders S, Kim V & Huber W. RNA-Seq workflow: gene-level exploratory
431 analysis and differential expression. *F1000Res* 2015; **4**:1070.

432 34 Piccaluga, PP, Agostinelli C, Califano A, Rossi M, Basso K, Zupo S *et al.* Gene
433 expression analysis of peripheral T cell lymphoma, unspecified, reveals distinct
434 profiles and new potential therapeutic targets. *J Clin Invest* 2007; **117**:823-834.

435 35 Agnelli, L, Mereu E, Pellegrino E, Limongi T, Kwee I, Bergaggio E *et al.* Identification
436 of a 3-gene model as a powerful diagnostic tool for the recognition of ALK-negative
437 anaplastic large-cell lymphoma. *Blood* 2012; **120**:1274-1281.

438 36 Wang, M, Zhang S, Chuang SS, Ashton-Key M, Ochoa E, Bolli N *et al.*
439 Angioimmunoblastic T cell lymphoma: novel molecular insights by mutation profiling.
440 *Oncotarget* 2017; **8**:17763-17770.

441 37 Inghirami, G, Chan WC, Pileri S & malignancies AxcG-dtmol. Peripheral T-cell and
442 NK cell lymphoproliferative disorders: cell of origin, clinical and pathological
443 implications. *Immunol Rev* 2015; **263**:124-159.

444 38 Reddy, A, Zhang J, Davis NS, Moffitt AB, Love CL, Waldrop A *et al.* Genetic and
445 Functional Drivers of Diffuse Large B Cell Lymphoma. *Cell* 2017; **171**:481-494 e415.

446 39 Scott, DW, Mottok A, Ennishi D, Wright GW, Farinha P, Ben-Neriah S *et al.*
447 Prognostic Significance of Diffuse Large B-Cell Lymphoma Cell of Origin Determined
448 by Digital Gene Expression in Formalin-Fixed Paraffin-Embedded Tissue Biopsies. *J*
449 *Clin Oncol* 2015; **33**:2848-2856.

450 40 Veldman-Jones, MH, Lai Z, Wappett M, Harbron CG, Barrett JC, Harrington EA *et al.*
451 Reproducible, Quantitative, and Flexible Molecular Subtyping of Clinical DLBCL
452 Samples Using the NanoString nCounter System. *Clin Cancer Res* 2015; **21**:2367-
453 2378.

454 41 Genovese, G, Kahler AK, Handsaker RE, Lindberg J, Rose SA, Bakhoum SF *et al.*
455 Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *N*
456 *Engl J Med* 2014; **371**:2477-2487.

457 42 Jaiswal, S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG *et al.* Age-
458 related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med* 2014;
459 **371**:2488-2498.

460

461